


STRONG ANOMALIES IN KAON DECAYS

S. TRINE

*Institut de Physique Théorique et Mathématique,
Université Catholique de Louvain, Chemin du Cyclotron, 2,
B-1348 Louvain-la-Neuve, Belgium*



We point to a potentially large contribution of the QCD trace anomaly to the isoscalar pion pair production in Kaon decays. This contribution arises from the long-distance evolution of a 7-dimensional quark-gluon operator generated by short-distance QCD corrections to the $\Delta S = 1$ Fermi Hamiltonian.

1 Introduction

In addition to the well-known anomaly in the global $U(1)$ axial symmetry, massless QCD also exhibits a breaking of the invariance under dilatations at the quantum level. This is known as the trace anomaly. While the axial anomaly is related to a pseudoscalar combination of two gluon fields, the trace anomaly is associated with a scalar combination of two such fields.

Their implications in the hadronic world are numerous. For example, in the quarkonium transitions $\Psi' \rightarrow J/\Psi\eta$ and $\Psi' \rightarrow J/\Psi\pi\pi$, it has been shown¹ that the gluon conversions into η and $\pi\pi$ are governed by the axial and trace anomalies respectively.

Similarly, the possibility of trace anomaly dominance has been investigated in weak K decays.² In this talk, we show how an enhancement of the isoscalar pion pair production due to trace anomaly effects may be induced by strong interaction corrections to the $\Delta S = 1$ Fermi Hamiltonian.

2 A trace anomaly contribution to the $\Delta I = 1/2$ rule in Kaon decays

The lowest order $\Delta S = 1$ quark-gluon (qG) effective Hamiltonian that contains the trace anomaly (TA) operator reads:

$$H_{TA}^{(qG)} \sim \bar{s}_R d_L G_{\mu\nu}^a G^{\mu\nu,a}, \quad (1)$$

where $G_{\mu\nu}^a$ denotes the gluon field strength tensor and the subscript $L(R)$ refers to left(right)-handed spinors.

This 7-dimensional operator is generated by leading order short-distance (SD) QCD corrections to the $\Delta S = 1$ Fermi Hamiltonian, at next-to-leading order in the momentum expansion.³ Yet the bulk of its contribution is expected to come from long-distance (LD) physics, as the anomaly is a non-perturbative phenomenon. Its effects on the $\Delta I = 1/2$ amplitudes could thus be larger than first expected from power counting arguments.

The LD evolution is computed in a truncated non-linear sigma model approach. The initial operator (1) is bosonized into U^{ds} at the confining scale. The unitary matrix U is related to the octet of light pseudoscalar fields π^a through $U = \exp(i\sqrt{2}\Pi/F_\pi)$, $\Pi = \sum_{a=1}^8 \lambda^a \pi^a / \sqrt{2}$. F_π is the neutral pion decay constant and λ^a are the Gell-Mann matrices with normalization $\text{tr} \lambda^a \lambda^b = 2\delta^{ab}$. Strong interaction corrections are modelled by the usual $O(p^2)$ non-linear Lagrangian $\mathcal{L}_{NL} = (F_\pi^2/4)\text{tr}(\partial_\mu U \partial^\mu U^\dagger) + r(F_\pi^2/4)\text{tr}(MU^\dagger + UM)$, where M is the real, diagonal, light quark mass matrix.

Using the background field method with a cut-off regularization, we obtain the following result at the one-loop level:^a

$$\begin{aligned} U^{ds} &\rightarrow (1+a)U^{ds} \\ &+ (b/2F_\pi^2)U^{ds}\text{tr}(\partial_\mu U \partial^\mu U^\dagger + r(MU^\dagger + UM)) \\ &+ (3b/2F_\pi^2)(\partial_\mu U \partial^\mu U^\dagger + r(MU^\dagger + UM))^{dq}U^{qs} + O(p^6) \end{aligned} \quad (2)$$

with $a = -3\Lambda^2/(4\pi F_\pi)^2$, $b = \ln(\Lambda^2/m_\pi^2)/32\pi^2$ and Λ is the ultra-violet cut-off.

We see that the lowest order non-linear realization of the TA operator, obtained by factorizing the quark from the gluon fields in (1), does indeed show up in the second line of Eq. (2).

3 Conclusion

We have indicated how a potentially important contribution of the QCD trace anomaly to the $\Delta I = 1/2$ Kaon decay amplitudes can arise from strong interaction corrections to the $\Delta S = 1$ Fermi Hamiltonian. A more quantitative analysis will require the investigation of the matching between the SD and LD evolutions. A previous study⁵ of $K \rightarrow \pi\pi$ decay amplitudes with gluons in the intermediate state (but without any reference to the trace anomaly) predicts an enhancement of $14 \pm 6\%$ using QCD sum rules. The consistency between the two methods would be worth checking.

Acknowledgments

This work has been done in collaboration with J.-M. Gérard.⁶ Financial supports from IISN and Interuniversity Attraction Pole P5/27 are also acknowledged.

References

1. M. Voloshin and V. Zakharov, *Phys. Rev. Lett.* **45**, 688 (1980);
V.A. Novikov and M.A. Shifman, *Z. Phys. C* **8**, 43 (1981).
2. J.-M. Gérard and J. Weyers, *Phys. Lett. B* **503**, 99 (2001).
3. A.A. Penin and A.A. Pivovarov, *Phys. Rev. D* **49**, 265 (1994).
4. T. Hambye *et al.*, *Phys. Rev. D* **58**, 014017 (1998).
5. A.A. Penin and A.A. Pivovarov, *Nuovo Cim. A* **107**, 1211 (1994).
6. J.-M. Gérard and S. Trine, in preparation.

^aNote that a similar computation has already been performed in another context.⁴